arXiv:1208.5220v1 [math.GR] 26 Aug 2012

OBSTRUCTIONS TO NONPOSITIVE CURVATURE FOR OPEN MANIFOLDS

IGOR BELEGRADEK

Abstract. We study algebraic conditions on a group G under which every properly discontinuous, isometric G-action on a Hadamard manifold has a G-invariant Busemann function. For such G we prove the following structure theorem: every open complete nonpositively curved Riemannian $K(G,1)$ manifold that is homotopy equivalent to a finite complex of codimension ≥ 3 is an open regular neighborhood of a subcomplex of the same codimension. In this setting we show that each tangential homotopy type contains infinitely many open $K(G,1)$ manifolds that admit no complete nonpositively curved metric even though their universal cover is the Euclidean space. A sample application is that an open contractible manifold W is homeomorphic to a Euclidean space if and only if $W \times S^1$ admits a complete Riemannian metric of nonpositive curvature.

1. Introduction

Throughout the paper unless stated otherwise all manifolds are smooth and connected, all metrics are complete Riemannian, and the phrase "nonpositive sectional curvature" is abbreviated to "$K \leq 0$".

A manifold is *covered by* $\mathbb{R}^n$ if its universal cover is diffeomorphic to $\mathbb{R}^n$. The Cartan-Hadamard theorem yields a basic obstruction to $K \leq 0$: any n-manifold with a metric of $K \leq 0$ is covered by $\mathbb{R}^n$.

Recall that an open $K(G,1)$ manifold covered by $\mathbb{R}^n$ exists if and only if G is a countable group of finite cohomological dimension. Countable groups of finite cohomological dimension form a class of torsion-free groups that is closed under subgroups, extensions, and amalgamated products.

A much deeper obstruction to $K \leq 0$ arises from Gromov's random, torsion-free, hyperbolic groups with strong fixed point properties: Each of the groups can be realized as the fundamental group of open manifold covered by $\mathbb{R}^n$ for large enough n, yet no such manifold is homotopy equivalent to a manifold of $K \leq 0$, see [Gro03] and [NS11, Theorem 1.1].

2000 Mathematics Subject classification. Primary 53C21, Secondary 57N15, 57S30. *Keywords:* nonpositive curvature, Busemann function, obstructions.

Even though the focus of this paper is on open manifolds, we next review known obstructions to $K \leq 0$ for closed manifolds. There exist closed manifolds that are covered by $\mathbb{R}^n$ and have one of the following properties: their fundamental groups are not CAT(0), see [Mes90, Dav08, Sap], they are homeomorphic but not diffeomorphic to manifolds of $K \leq 0$ [Ont03], they admit CAT(0) metrics but are not homotopy equivalent to closed manifold of $K \leq 0$ [DJL12].

A manifold is *covered by* $\mathbb{R} \times \mathbb{R}^{n-1}$ if it is diffeomorphic to the product of $\mathbb{R}$ and a manifold covered by $\mathbb{R}^{n-1}$. In the paper we describe a sizable class of groups such that for each group G in the class

(i) every $K(G,1)$ manifold of $K \leq 0$ is covered by $\mathbb{R} \times \mathbb{R}^{n-1}$;

(ii) there are $K(G,1)$ manifolds that are covered by $\mathbb{R}^n$ but not homeomorphic to manifolds covered by $\mathbb{R} \times \mathbb{R}^{n-1}$.

This yields a homeomorphism obstruction to $K \leq 0$ for if G satisfies (i)-(ii), then no manifold in (ii) is homeomorphic to a manifold of $K \leq 0$.

Manifolds covered by $\mathbb{R} \times \mathbb{R}^{n-1}$ arise in the context of nonpositive curvature, namely, consider a manifold of $K \leq 0$ written as X/G where G is the deck transformations group acting isometrically on the universal cover X. If G *fixes a Busemann function* (i.e. G fixes a point at infinity and stabilizes each horosphere centered at the point), then X/G is covered by $\mathbb{R} \times \mathbb{R}^{n-1}$. More precisely, any Busemann function is a C^2-submersion [HIH77], so it defines a C^1 diffeomorphism of X and $\mathbb{R} \times \mathbb{R}^{n-1}$, and if the Busemann function is G-invariant, the C^1-diffeomorphism descends to quotients, which are therefore C^∞-diffeomorphic.

The property "G fixes a Busemann function" can be forced by purely algebraic assumptions on G summarized in Sections 2-3, where we also describe various classes of groups with this property some of which are listed below.

Example 1.1. Any of the following conditions implies that each isometric, properly discontinuous G-action on a Hadamard manifold X fixes a Busemann function:

(1) $G = \pi_1(E)$ where E is a closed aspherical manifold whose finitely-sheeted cover is the total space of an oriented circle bundle with nonzero rational Euler class. This applies when E is an infranilmanifold of nilpotency degree ≥ 2, or a 3-manifold modelled on $\widetilde{SL}_2(\mathbb{R})$.
(2) $G = \pi_1(E) \times H$ where E is as in (1) and H is any group.
(3) $G = \mathbb{Z}^n \rtimes H$ where $n \geq 3$ and H be a finite index subgroup of $GL_n(\mathbb{Z})$, and the semidirect product is defined via the standard H-action on $\mathbb{Z}^n$.
(4) G is any group with infinite center and finite abelianization, e.g. the preimage of any lattice under the universal cover $\widetilde{Sp}_{2n}(\mathbb{R}) \to Sp_{2n}(\mathbb{R})$, $n \geq 2$.

(5) G is the amalgamated product along $\mathbb{Z}^n$ of the group $\mathbb{Z}^n \rtimes H$ as in (4) and any group K that contains $\mathbb{Z}^n$ as a normal subgroup.
(6) G has infinitely generated center, e.g. the product of $(\mathbb{Q}, +)$ and any group.

In Section 4 we show that if the Hadamard manifold X is *visibility* (i.e. any two distinct points at infinity are the endpoints of a geodesic in X), then the class of groups that fix Busemann functions on X gets even larger, e.g. it contains any nontrivial product $G = G_1 \times G_2$ such that there is a $K(G,1)$ manifold.

An essential tool in understanding manifolds covered by $\mathbb{R} \times \mathbb{R}^{n-1}$ is the recent result of Guilbault [Gui07]: if an open manifold W of dimension ≥ 5 is homotopy equivalent to a finite complex, then $\mathbb{R} \times W$ is diffeomorphic to the interior of a compact manifold. As an addendum to [Gui07] we show:

Theorem 1.2. *Let W be an open $(n-1)$-manifold with $n \geq 5$ that is homotopy equivalent to a finite complex of dimension $k \leq n-3$. Then $\mathbb{R} \times W$ is diffeomorphic to the interior of a regular neighborhood of a k-dimensional finite subcomplex.*

In addition to fixing a Busemann function there are other properties that can force any open $K(G,1)$ manifold of $K \leq 0$ to be the interior of a regular neighborhood of a finite subcomplex.

Example 1.3. Let G be the fundamental group of a closed flat manifold such that either $G \cong \mathbb{Z}$ or G has finite abelianization, see [Szc85] for examples. Then for any $K(G,1)$ manifold with $K \leq 0$, the G-action on the universal cover X either fixes a Busemann function or acts cocompactly on a totally geodesic stabilizes a flat F where it acts cocompactly; in the latter case X/G is diffeomorphic to the normal bundle of the closed flat submanifold F/G, see Section 2.

Example 1.4. Let G be an irreducible torsion-free uniform lattice in a connected higher rank semisimple Lie group with finite center and no compact factor. Then harmonic maps superrigidity [Duc, Theorem 1.2] implies that for any $K(G,1)$ manifold with $K \leq 0$, the G-action on the universal cover X either acts cocompactly on a totally geodesic subspace F, or fixes a point at infinity in which case it also fixes the corresponding Busemann function as G^{ab} is finite (see Remark 3.3).

Theorem 1.2 and Example 1.3 yield an attractive characterization of $\mathbb{R}^n$:

Corollary 1.5. *An open contractible n-manifold W is homeomorphic to $\mathbb{R}^n$ if and only if $W \times S^1$ admits a metric of $K \leq 0$.*

If $n = 4$, then "homeomorphic" in Corollary 1.5 cannot be replaced with "diffeomorphic" because if W is an exotic $\mathbb{R}^4$, then $W \times S^1$ is diffeomorphic to $\mathbb{R}^4 \times S^1$ [Sie69, Theorem 2.2].

Since there are only countably many diffeomorphism classes of compact manifolds, there are only countably many possibilities for open regular neighborhoods. Various classification results for manifolds diffeomorphic to the interiors of regular neighborhoods can be found in [Hae61, RS68, Sie69, LS69] and [Wal99, Corollary 11.3.4]. In the stable range, i.e. when the subcomplex has dimension $< \frac{n}{2}$, the interior of a regular neighborhood of dimension n is determined up to diffeomorphism by its tangential homotopy type [LS69].

Corollary 1.6. *Let G be a group as in* Examples 1.1, 1.3, 1.4. *If V is an open n-manifold that is homotopy equivalent to a finite $K(G,1)$ complex of dimension $< \frac{n}{2}$, where $n \geq 5$, then the tangential homotopy type of V contains at most one manifold of $K \leq 0$.*

In metastable range one knows [Hae61, Sie69] that if a regular neighborhood is homotopy equivalent to a closed manifold, then its interior is the total space of a vector bundle over that manifold. The metastable range result together with Theorem 1.2 implies:

Corollary 1.7. *Let B be a closed aspherical k-manifold as in* Example 1.1(1) *or* Examples 1.3-1.4, *and let V be an open n-manifold of $K \leq 0$ that is homotopy equivalent to B. If $2n > 3k + 2$ and $n > 4$, then V is diffeomorphic to the total space of a vector bundle over B.*

We now turn to methods of producing manifolds as in (ii). A trivial method is to consider a $K(G,1)$ manifold that has the minimal dimension among all $K(G,1)$ manifolds that are covered by $\mathbb{R}^n$, which yields:

Proposition 1.8. *Any aspherical manifold is homotopy equivalent to a manifold covered by $\mathbb{R}^n$ but not covered by $\mathbb{R} \times \mathbb{R}^{n-1}$.*

Combining the proposition Example 1.1(6) we get:

Corollary 1.9. *If G has infinitely generated center, then any $K(G,1)$ manifold is homotopy equivalent to a manifold that admits no metric of $K \leq 0$.*

This trivial method always works, yet it is non-constructive for it is not easy to decide whether a specific open manifold has the minimal dimension in the above sense, see [BKK02, BF02, Yoo04, Des06].

On the other hand, if G has a finite $K(G,1)$ complex, we produce explicit manifolds satisfying (ii):

Theorem 1.10. *Let V be an open aspherical n-manifold that is homotopy equivalent to a finite complex of dimension $\le n-3$, where $n \ge 5$. Then the tangential homotopy type of V contains countably many manifolds such that each of them is covered by $\mathbb{R}^n$ and is not homeomorphic to the interior of a regular neighborhood of a finite subcomplex of codimension ≥ 3, and in particular not covered by $\mathbb{R} \times \mathbb{R}^{n-1}$.*

It should be possible to strengthen the conclusion "countably many" in Theorem 1.10 to "a continuum of", but we only prove this under a technical assumption which holds in many cases of interest, including when $\pi_1(V)$ is a Poincaré duality group, or when $\pi_1(V)$ does not contain a nonabelian free subgroup, see Section 6.

The structure of the paper is as follows: Sections 2, 3, 4 discuss when a group fixes a Busemann function, and in particular, justify Examples 1.1, 1.3, 1.4. In Section 5 we study manifolds with $\mathbb{R}$-factors, and prove Theorem 1.2 and Corollary 1.5. In Section 6 we construct manifolds without $\mathbb{R}$-factors, and prove Theorem 1.10 as well as its harder uncountable version. The latter requires a certain family of homology 3-spheres described in Appendix B. Throughout the paper we make implicit use of various finiteness results for groups, which are recalled in Appendix A.

Acknowledgments. The author is grateful for NSF support (DMS-1105045) and to MathOverflow for stimulating work environment. Thanks are due to Karel Dekimpe for Examples 3.8 and 3.10, Misha Kapovich for Example 3.9, Andy Putman and Wilberd van der Kallen for help with Proposition 3.12.

2. Parabolics in the center and invariant Busemann functions

Background references for nonpositive curvature are [BGS85, Ebe96, BH99]. In this section G is a discrete isometry group of a Hadamard manifold X.

Lemma 2.1. *If W is a closed convex subset of X, then any abelian discrete subgroup $A \le Iso(W)$ that contains no parabolic isometries is finitely generated of rank $\le \dim(W)$.*

Proof. By [BGS85, lemma 7.1(1)] the set $\bigcap_{g\in A} \mathrm{Min}(g)$ is a closed A-invariant convex subset of M which splits as $C \times \mathbb{R}^k$, where A acts trivially on the C-factor, and by translations on $\mathbb{R}^k$. Since A is discrete, it acts freely on $\mathbb{R}^k$ so $A \cong \mathbb{Z}^m$ for $m \le k$. □

Lemma 2.2. *G preserves a Busemann function if it has a finite index subgroup G_0 whose center $Z(G_0)$ contains a parabolic isometry.*

Proof. This result is implicit in [BGS85, Lemma 7.3, 7.8], but for completeness we record a proof below. Let $z \in Z(G_0)$ be a parabolic isometry, and $g_1, \dots g_k$ are right coset representatives of G_0 in G, then the function $x \to \sum_i d(g_i^{-1} z g_i x, x)$ is convex, and G-invariant because $d(g_i^{-1} z g_i x, x)$ is independent of a coset representative in $G_0 g_i$ as $z \in Z(G_0)$. Since $x \to d(zx, x)$ does not assume its infimum, neither does the above convex function. Then a limiting process outlined in [BGS85, Lemma 3.9], and fully explained in [BH99, Lemma II.8.26], gives rise via Arzela-Ascoli theorem to a G-invariant Busemann function. Note that this Busemann function is not explicit and there seem to be no way to find a G-invariant Busemann function associated with a given point of $X(\infty)$. □

Theorem 2.3. *Let G be a group with subgroups H, G_0 such that their centers $Z(H)$, $Z(G_0)$ are infinite, $Z(H) \subseteq Z(G_0)$, the index of G_0 in G is finite, and one of the following conditions hold:*
(1) *$Z(H)$ is not finitely generated;*
(2) *any homomorphism $H \to \mathbb{R}$ is trivial.*
(3) *H is finitely generated, and $Z(H)$ contains a free abelian subgroup that is not a direct factor of any finite index subgroup of H.*
Then any effective, properly discontinuous, isometric G-action on a Hadamard manifold fixes a Busemann function.

The reader may want to first think through the case when $H = G_0 = G$, and then go on to observe that if Theorem 2.3 applies to H, G_0, G, then it also applies to H, $G_0 \times K$, $G \times K$ for any group K.

Proof. The result follows by Lemma 2.2 once we show that $Z(G_0)$ contains a parabolic isometry. Since $Z(H) \subseteq Z(G_0)$ it suffices to find a parabolic element in $Z(H)$. Suppose no such element exists. Then $Z(H)$ is finitely generated by Lemma 2.1, which implies (1).

To prove (2) note that if H has no nontrivial homomophism into $\mathbb{R}$, then by [BH99, Remark II.6.13] $Z(H)$ contains no axial isometries, so it is a finitely generated, abelian, and periodic, and hence finite which contradicts the assumptions.

Finally, (3) is deduced as follows. Since $Z(H)$ is finitely generated and infinite, it has a free abelian subgroup A of positive rank. Under the assumption that H is finitely generated, [BH99, Theorem II.7.1] implies that H has a finite index subgroup that contains A as a direct factor. □

Example 2.4. Theorem 2.3(1) applies, e.g. to any infinitely generated, torsion-free, countable abelian group of finite rank, such as $(\mathbb{Q}, +)$, where finiteness of rank ensures finiteness of cohomological dimension [Bie81, Theorem 7.10].

Question 2.5. *Is there a group of type F with infinitely generated center?*

Remark 2.6. If H is a group with infinitely generated center, then of course, H cannot be embedded into hyperbolic or $\mathrm{CAT}(0)$ groups because their abelian subgroup are finitely generated. If in addition H has type *F*, or even of type *FP*, then H cannot be linear over a field of characteristic zero [AS82, Corollary 5], and cannot be elementary amenable, as we explain below for completeness.

Proposition 2.7. *Every elementary amenable group of type FP has finitely generated center.*

Proof. Since G has type *FP*, its cohomological dimension is finite. Any elementary amenable of finite cohomological dimension is virtually solvable [HL92] and of course torsion-free. As G has type *FP*, so does the solvable finite index subgroup G_0 of G [Bro94, Theorem VIII.6.6]. Solvable groups of type *FP*, are constructable [Kro86] and hence quotients of G_0 are finitely presented [BB76, Theorem 4]. So $Z(G_0)$ is finitely generated, for if the quotient of a finitely generated group by the center is finitely presented, then the center is finitely generated [Bau93, Corollary 1, page 54]. Hence the subgroup $Z(G) \cap G_0$ of $Z(G_0)$ is finitely generated, and therefore the same holds for $Z(G)$. □

Example 2.8. Here is an example of a group of type *F* with infinite center and finite abelianization (to which Theorem 2.3(2) applies). Let G be the preimage of a lattice $\Gamma \leq Sp_{2n}(\mathbb{R})$ under the universal cover, then $Z(G)$ is infinite, and G has Kazhdan property (T) when $n \geq 2$ [BdlHV08, Example 1.7.13], and hence finite abelianization. If Γ is torsion-free, then it has type *F* [BS73], and hence so does G, see Appendix A. Note that Theorem 2.3(3) also applies to G because its finite index subgroups have property (T) and hence finite abelianization which implies that they cannot have $\mathbb{Z}$ as a direct factor.

Theorem 2.3(3) also applies when G is virtually nilpotent, finitely generated, and not virtually abelian, namely, we let $H = G_0$ be a nilpotent torsion-free subgroup of finite index, and use the following.

Proposition 2.9. *If H is a torsion-free, nonabelian, nilpotent group H, then $Z(H)$ is not a virtual direct factor of H.*

Proof. Arguing by contradiction suppose H_0 is a finite index subgroup of H that contains $Z(H)$ as a direct factor, i.e. there is $K \trianglelefteq H_0$ with $H = KZ(H)$ and $K \bigcap Z(H) = 1$. Then $Z(H_0) = Z(K)Z(H)$ where $Z(K) \bigcap Z(H) = 1$. A key observation is that $Z(H_0) \subseteq Z(H)$: if we take $z \in Z(H_0)$ and $g \in H$, and pick k with $g^k \in H_0$, then $g^k = (zgz^{-1})^k$ but in torsion-free nilpotent groups roots are unique [KM79, Theorem 16.2.8], so $g = zgz^{-1}$, and hence $z \in Z(H)$. It follows that $Z(K)$ is trivial, so since K is nilpotent we conclude that K is trivial, i.e. $H_0 = Z(H)$. Now if $g, h \in H$, then $h^k \in H_0$ so $gh^kg^{-1} = h^k$ and again uniqueness of roots implies $ghg^{-1} = h$ so that H is abelian. □

It is a well-known fact that the group H in Proposition 2.9 can be realized as the fundamental group of the circle bundle over a closed nilmanifold, and noncommutativity of H allows to choose the bundle so that its rational Euler class is nonzero. Thus Proposition 2.9 is implied by the proposition below.

Proposition 2.10. *If B is a closed aspherical manifold, and $E \to B$ is a principal circle bundle with nonzero rational Euler class, then the fiber circle is a nontrivial element s in the center of $\pi_1(E)$ that is not a virtual direct factor.*

Proof. The circle action on E lifts to any cover $\hat{E}$ whose fundamental group contains s [Bre72, Theorem I.9.3]. Therefore, the covering $\hat{E} \to E$ descends to a covering of base spaces $\hat{B} \to B$. It follows that $\hat{E} \to \hat{B}$ is a pullback of $E \to B$ via $\hat{B} \to B$, but then $\hat{E} \to \hat{B}$ has nonzero rational Euler class because finite covers induce injective maps on rational cohomology (due to existence of a transfer). Finally, if s were a direct factor in $\pi_1(\hat{E})$, then $\hat{E} \to \hat{B}$ would have a section, so the Euler class would be zero. □

Remark 2.11. Theorem 2.3(3) also applies to the fundamental group of any closed Seifert fibered space modelled on $\widetilde{SL}_2(\mathbb{R})$, because it has a finite index subgroup to which Proposition 2.10 applies.

Further examples of groups that satisfy Theorem 2.3(3) by amalgamating one such example with any group along a common central subgroup, which uses the following fact.

Fact 2.12. *Let $G := G_1 *_A G_2$ where A lies in the center of G_1, G_2. If $B \leq A$ is a virtual direct factor of G, then B is a virtual direct factor of G_1 and G_2.*

Proof. Note that $B \leq Z(G)$. If K is a finite index subgroup of G of the form BL, where L is normal in K and $B \cap L = \{1\}$, then $K \cap G_i$ is a finite index subgroup of G_i of the form $B(L \cap G_i)$ where $L \cap G_i$ is normal in $K \cap G_i$ and $B \cap (L \cap G_i) = \{1\}$, so B is a virtual direct factor of G_i. □

3. GROUNDED GROUPS AND INVARIANT BUSEMANN FUNCTIONS

In this section G is a discrete isometry group of a Hadamard manifold X.

If G fixes a point at infinity, then G permutes Busemann functions (or equivalently, permutes horospheres) associated with the point.

Example 3.1. An axial isometry $z \to 2z$ of the upper half permutes horospheres centered at 0, as well as horospheres centered at ∞.

Fact 3.2. *If G fixes a point $[b]$ at infinity represented by a Busemann function b, then the map $g \to b(gx) - b(x)$ is a group homomorphism $G \to (\mathbb{R}, +)$ that depends only on $[b]$, and not on x or b.*

Proof. This is immediate from the cocycle property of the Busemann function, e.g. see the discussion after [CM09, Proposition 3.11] where the above homomorphism was called the *Busemann character.* ☐

Remark 3.3. Thus if $G \le \mathrm{Iso}(X)$ has no nontrivial homomorphism into $\mathbb{R}$, then G preserves a Busemann function if and only if G fixes a point at infinity.

To further enlarge the class of groups that preserve Busemann functions we discuss, following Schroeder in [BGS85, Appendix 3] and Eberlein [Ebe96, Section 4.4], groups which we call *grounded* (and which were called *admissible* in [Ebe96]). Namely, equip the ideal boundary $X(\infty)$ of a Hadamard manifold X with the Tits metric, which is $CAT(1)$ and complete [BH99, Theorem II.9.20], and call a subgroup $G \le \mathrm{Iso}(X)$ *grounded* if its fixed point set at infinity $X_G(\infty)$ is a nonempty subset of (intrinsic) radius $\le \frac{\pi}{2}$, i.e. $X_G(\infty)$ lies in the closed $\frac{\pi}{2}$-ball centered at a point of $X_G(\infty)$.

Example 3.4. There are discrete cocompact subgroups of $\mathrm{Iso}(\mathbb{R}^n)$ with finite abelianizations, such as $(\mathbb{Z}_2 * \mathbb{Z}_2)^n$, or see [Szc85] for torsion-free examples. No such group can fix a point at infinity of $\mathbb{R}^n$ because no lattice fixes a Busemann function. By contrast, any group of translations, such as the standard $\mathbb{Z}^n$, fixes the ideal boundary of $\mathbb{R}^n$ pointwise, yet it is not grounded as its Tits boundary is isometric to the standard unit sphere, which has radius π.

Theorem 3.5. *A discrete, infinite isometry group G of a Hadamard manifold is grounded if one of the following is true:*
(1) *G is as in* Theorem 2.3.
(2) *G has a grounded normal subgroup.*
(3) *G is virtually solvable and not virtually-$\mathbb{Z}^k$ for any k.*
(4) *G is the union of a nested sequence of grounded subgroups.*
(5) *a normal abelian subgroup of G contains an infinite G-conjugacy class.*

Proof. Schroeder proved in [BGS85, Appendix 3] the key fact that an abelian group that contains a parabolic is grounded, see also [Ebe96, Section 4.4]. These references also contain a proof of part (2) which we outline below. If N is a grounded normal subgroup of G, then G stabilizes $X_N(\infty)$, and hence fixes its center of gravity, which proves that $X_G(\infty)$ is nonempty. It is clear that $X_G(\infty) \subseteq X_N(\infty)$ and that $X_G(\infty)$ is closed in the cone topology on $X(\infty)$. Since the identity map of $X(\infty)$ from the Tits metric topology to the cone the topology is continuous, $X_G(\infty)$ is closed in the Tits metric topology. Finally, convexity of $X_G(\infty)$ is implied by the fact that the $\frac{\pi}{2}$-balls are uniquely geodesic, so if an element of G fixes the endpoints of a geodesic segment, it fixes the segment pointwise.

To prove (4) let $G = \bigcup_i G_i$ where $G_1 \le \dots \le G_i \le \dots$ is a countable family of grounded subgroups of $\mathrm{Iso}(X)$ (since G is countable, it is not the union of uncountable family of distinct groups). Note that $X_G(\infty) = \bigcap_i X_{G_i}(\infty)$. By assumptions $X_{G_i}(\infty)$ is a nested family of closed convex subsets of radius $\le \frac{\pi}{2}$. It is shown by Caprace-Lytchak [CL10, Lemma 5.2] that any such family must have a nonempty intersection provided the ambient metric space is finite dimensional, which follows from [CL10, Proposition 2.1] because X is finite dimensional. Thus $X_G(\infty)$ is a nonempty, closed, convex subset of radius $\le \frac{\pi}{2}$, so G is grounded.

Now we are ready to prove (1). The conclusion would be immediate if G_0 were normal in G because by the proof of Theorem 2.3 $Z(G_0)$ contains a parabolic element, so G_0, and hence G is grounded by the above-mentioned result of Schroeder. In general, since G is countable, we can realize G as the union of a nested sequence of finitely generated subgroups $\{H_i\}$ where $H_i \subset H_{i+1}$ for all i. We may also assume that H_1 contains a parabolic element $z \in Z(G_0)$. The subgroup $N_i := \bigcap_{g\in H_i} g(H_i \cap G_0)g^{-1}$ is normal in H_i, and moreover, the index of N_i in H_i is finite because H_i is a finitely generated, and the index of $H_i \cap G_0$ is finite. Hence N_i contains a power of z, which is a parabolic element in $Z(N_i)$. It follows that N_i is grounded, and hence so is H_i. Now (4) implies that G is grounded as the union of H_i's.

To prove (3) we first recall a result of Burger-Schroeder [BS87] that an amenable subgroup of $\mathrm{Iso}(X)$ either stabilizes a flat, or fixes a point of $X(\infty)$. In the former case the subgroup is virtually-$\mathbb{Z}^k$, which is excluded by assumption. (If a discrete subgroup $G \le \mathrm{Iso}(X)$ stabilizes a m-dimensional flat, then restricting the G-action to the flat gives a homomorphism $G \to \mathrm{Iso}(\mathbb{R}^m)$ with finite kernel and discrete image. By Bieberbach's theorem the image of G stabilizes an k-dimensional flat in $\mathbb{R}^k$, which implies that G is virtually-$\mathbb{Z}^k$, see e.g [Wil00, Theorem 2.1]).

Thus we may assume that $X_G(\infty)$ is nonempty. By Lemma 2.1 abelian subgroups of G are finitely generated, so G is virtually polycyclic as proved by Mal'cev [Mal51]. A polycyclic subgroup of $\mathrm{Iso}(X)$ that contains no parabolics must be virtually abelian [BH99, Theorem II.7.16]. Thus G contains a parabolic element. If P is the cyclic subgroup of G generated by a parabolic element, then $X_G(\infty) \subseteq X_P(\infty)$, where $X_P(\infty)$ has radius $\le \frac{\pi}{2}$ because P is grounded.

To prove (5) consider an normal abelian subgroup A of G that contains an infinite G-conjugacy class. Then A contains a parabolic element [BH99, Lemma II.7.17(2)], which implies that A, and hence G are grounded. □

Corollary 3.6. *Let G be a finitely generated, torsion-free, discrete subgroup of Iso(X). If G has a nontrivial, normal, elementary amenable subgroup, then either G is grounded, or G has a nontrivial, finitely generated, abelian, normal subgroup that is a virtual direct factor of G.*

Proof. Since G is torsion-free and discrete, it has finite cohomological dimension, so by [HL92, Corollary 2] G has a nontrivial, abelian, normal subgroup A. If A contains a parabolic element, then G is grounded. Otherwise, by Lemma 2.1 A is isomorphic to some $\mathbb{Z}^k$, $k > 0$, and then [BH99, Theorem II.7.1(5)] implies that A is a virtual direct factor of G. □

Remark 3.7. If G is an amenable, discrete, torsion-free isometry group of X that is not virtually-$\mathbb{Z}^k$ for any k, then G fixes a point at infinity [BS87], yet I do not know whether it is always grounded. A counterexample, if it exists, would have to be an amenable group of finite cohomological dimension that is not elementary amenable, for if G were elementary amenable, it would be virtually solvable [HL92], and hence grounded by Theorem 3.5(3).

In view of Remark 3.3 one wants to produce grounded groups with no nontrivial homomorphisms into $\mathbb{R}$.

Example 3.8. A complete list of 3-dimensional closed infranilmanifolds, as found in [DIKL95], contains an infinite family of manifolds whose fundamental group has finite abelianization. Namely, in the family 2 on [DIKL95, page 156], also listed on [Dek96, page 160], let k_1 be a positive even integer, and set $k_2 = 0 = k_3$, $k_4 = 1$. Taking products yields examples in each dimension divisible by 3. This example does not really illustrate the power of Theorem 3.5(3) because it is already covered by Theorem 2.3, yet it can be used as a building block in other examples below. For that purpose we note that any group in the above family surjects onto the infinite dihedral group.

The following virtually solvable group has finite abelianization, finite cohomological dimension, and is not virtually polycyclic, so it satisfies assumptions of Theorem 3.5(3).

Example 3.9. (Misha Kapovich) Let H be a virtually polycyclic group of finite cohomological dimension such that H^{ab} is a finite group that has $\mathbb{Z}_m$ among its cyclic factors for even m (e.g. we could take H to be the fundamental group of Hantsche-Wendt's flat manifold, or any group in Example 3.8). Set $B := \mathbb{Q} \rtimes \mathbb{Z}$, where $\mathbb{Z}$ acts on $\mathbb{Q}$ by $q \to -q$ via the surjection onto $\mathbb{Z}_2$. Set $G := B \rtimes H$, where H acts on B so that the action on $\mathbb{Q}$ is trivial and the action on $\mathbb{Z}$ is by $m \to -m$ (which is indeed an action as $\mathbb{Z}$ is abelian). By construction G^{ab} is finite, and G has finite cohomological dimension because its semidirect factors have this property, e.g. $\mathbb{Q}$ is locally cyclic. Finally, G

is not virtually polycyclic as it contains $\mathbb{Q}$, and abelian subgroups of virtually polycyclic groups are finitely generated.

In the above example G is not of type F (because it has $\mathbb{Q}$ as a virtual direct factor). The following is an example of a torsion-free polycyclic group with finite abelianization which is not virtually nilpotent, and like every virtually polycyclic group, it is the fundamental group of a closed aspherical manifold [AJ76].

Example 3.10. (Karel Dekimpe) Let H be any virtually polycyclic group with finite abelianization such that H surjects onto the infinity dihedral group $D_\infty = \langle a, b | b^2 = 1, aba = b\rangle$. Let

$$\alpha = \begin{bmatrix} Q & 0 \\ 0 & Q^{-1} \end{bmatrix} \qquad\qquad \beta = \begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix}$$

be matrices on $GL_{2n}(\mathbb{Z})$, where $Q \in GL_n(\mathbb{Z})$. Since $\beta^2 = I_{2n}$ and $\alpha\beta\alpha = \beta$, the map that sends a, b to α, β, respectively, extends to a homomorphism $D_\infty \to GL_{2n}(\mathbb{Z})$. It remains to choose Q so that the corresponding semidirect product $G := \mathbb{Z}^4 \rtimes H$ has finite abelianization, and is not virtually nilpotent. The abelianization of the semidirect product $A \rtimes B$ is $(A^{\mathrm{ab}})_B \times B^{\mathrm{ab}}$, where $(A^{\mathrm{ab}})_B$ is the coinvariants for the B-action on A^{ab}, i.e. the quotient of A by the subgroup generated by the set $\{z - \bar{\phi}_b(z) | z \in A^{\mathrm{ab}}, b \in B\}$, where $\bar{\phi}$ is ϕ followed by the projection $A \to A^{\mathrm{ab}}$. Take $Q \in SL_2(\mathbb{R})$ with $\mathrm{trace}(Q) \neq 2$ so that $A - I_2$ has nonzero determinant. By considering where α, β map the standard generators of $\mathbb{Z}^4$ one gets finiteness of the coivariants. To show that G is not virtually nilpotent, we assume in addition that $|\mathrm{trace}(Q)| > 2$, and consider the subgroup $H_0 \leq H$ that is the preimage of the subgroup $\langle a\rangle \leq D_\infty$, and note the $\mathbb{Z}^2$-summand of $\mathbb{Z}^4$ spanned by the first two basis vectors is H_0-invariant. Thus G has a subgroup $\mathbb{Z}^2 \rtimes H_0$, which surjects onto $\mathbb{Z}^2 \rtimes \langle a\rangle$. With our choice of Q the latter group is a lattice in SOL, a 3-dimensional solvable Lie group. Since lattices in SOL are not virtually nilpotent, neither is G, as being virtually nilpotent is inherited by subgroups and quotients.

The following gives a large family of grounded groups of the form $\mathbb{Z}^k \rtimes K$, which have finite abelianization and type F if K does. This applies e.g. if K is a torsion-free lattice in $GL_k(\mathbb{Z})$ where $k \geq 3$ and the semidirect product is defined via its standard action on $\mathbb{Z}^k$.

Proposition 3.11. Theorem 3.5(5) *applies to every semidirect product* $G = \mathbb{Z}^k \rtimes K$ *defined by a homomorphism* $\phi\colon K \to GL_k(\mathbb{Z})$ *whose image contains an infinite order element.*

Proof. If $h \in H$ is mapped to an infinite order element ϕ_h, then there is $v \in \mathbb{Z}^k$ such that $\phi_{h^m}(v) \neq v$ for $m \neq 0$ (e.g. v can be chosen as a vector of the standard basis $\{e_i\}$, for otherwise for each i we have $\phi_{h^{l_i}}(e_i) = e_i$ for some l_i, so if $l = l_1 \dots l_k$, then by linearity ϕ_{h^l} is the identity, contradicting the infinite order assumption). Thus $h^m v h^{-m} = \phi_{h^m}(v) \neq v$ in G for all $m \neq 0$ so the conjugacy class of a intersects $\mathbb{Z}^k$ along an infinite subset, and Theorem 3.5(5) applies. ☐

Proposition 3.12. *In the notations of* Proposition 3.11 *if $\phi(K)$ is a finite index subgroup of $GL_k(\mathbb{Z})$, and if K^{ab} is finite, then G^{ab} is finite. In particular, G^{ab} is finite if ϕ is injective and $n \geq 3$.*

Proof. As in Example 3.10, the abelianization of $A \rtimes B$ is $(A^{\mathrm{ab}})_B \times B^{\mathrm{ab}}$. Finiteness of $(\mathbb{Z}^k)_K$ follows via the elementary computation below shown to me by Wilberd van der Kallen. For $i \neq j$, let $E^m_{ij} \in GL_k(\mathbb{Z})$ be the matrix with 1's on the diagonal, m as ij-entry, and zero elsewhere. For every m, and any basis vector $e_l \in \mathbb{Z}^k$ one can be easily find $v \in \mathbb{Z}^k$ and E^m_{ij} such that $me_i = v - E^m_{ij}(v)$. Thus it remains to find m such that $\phi(K)$ contains every matrix E^m_{ij}. Let C be a finite index normal subgroup of $GL_k(\mathbb{Z})$ that lies in $\phi(K)$. Since the mth power of E^1_{ij} equals E^m_{ij}, it suffices to pick m divisible by the index of C in $GL_k(\mathbb{Z})$.

An elegant way to prove finiteness of $(\mathbb{Z}^k)_K$ was explained to me by Andy Putman: One needs to show that $(\mathbb{R}^k)_K = 0$, and since K is a lattice in $SL_k(\mathbb{R})$, the Borel density theorem shows that $\mathbb{R}^k$ is an irreducible K-module, so the submodule generated by $\{v - \phi_g(v) | v \in \mathbb{R}^k,\ g \in K\}$ cannot be proper.

Finally, if ϕ is injective and $n \geq 3$, then K has Kazhdan property (T), and hence K^{ab} is finite. ☐

Remark 3.13. The property of having finite abelianization is inherited by amalgamated products and extensions: the former is obvious and the latter follows from right exactness of the abelianization functor.

Example 3.14. Suppose A is a normal abelian subgroup of groups H and F, so that A is also normal in $G := H *_A F$. Suppose H and A satisfy an algebraic condition that force the existence of a parabolic in A for every isometric, properly discontinuous H-action on a Hadamard manifold, which happens e.g. if H is as Theorem 2.3, or if $H = \mathbb{Z}^k \rtimes K$ as in Proposition 3.12. Then G is grounded, and the above discussion allows to construct many such groups that have finite abelianization and type F.

4. INVARIANT BUSEMANN FUNCTIONS ON VISIBILITY MANIFOLDS

Grounded groups are much easier to find among isometry groups of visibility manifolds. The main geometric ingredient is supplied by the following.

Lemma 4.1. *If X is visibility, G is not virtually cyclic, and $X_G(\infty)$ is nonempty, then G fixes a Busemann function, G contains no axial isometry, $X_G(\infty)$ is a single point, and in particular G is grounded.*

Proof. If $X_G(\infty)$ contains two distinct points, then these points can be joined by a geodesic as X is visibility. The union of all geodesics that join the points splits as $C \times \mathbb{R}$, where C is a compact convex subset. There exists an element of G acts as a nontrivial translation on the $\mathbb{R}$ factor (for if G-action on the $\mathbb{R}$-factor were trivial, compactness of C and proper discontinuity of the action on $C \times \mathbb{R}$ would imply that finiteness of the group G). Thus G-action on $C \times \mathbb{R}$ is cocompact, so G is two ended and hence virtually-$\mathbb{Z}$, contradicting the assumptions. Thus $X_G(\infty)$ is a single point, to be denoted β.

First we deal with the case when G is periodic. Any elliptic element $g \in G$ pointwise fixes the ray that connects β with any point of X fixed by g, and hence g preserves the Busemann function for that ray, and hence every Busemann function for β as they differ by a constant.

A key property of a parabolic isometry of a visibility manifold is that is has one fixed point at infinity, and similarly, any axial isometry fixes precisely two points at infinity [BGS85, Lemma 6.8].

It follows from [KN04, Proposition 4, page 8] that the fixed point sets of any two infinite order isometries are either disjoint or equal. Since any element of G fixes β, all infinite order isometries in G have the same fixed point set.

Suppose G contains an axial isometry and no parabolic isometries. Let $\alpha \in X(\infty) - \{\beta\}$ such that each axial isometry in G fixes α and β. If $e \in G$ is elliptic and $h \in G$ is axial, then ehe^{-1} is axial with fixed points $e(\beta) = \beta$ and $e(\alpha)$, so $e(\alpha) = \alpha$. This contradicts that above conclusion that $X_G(\infty) = \{\beta\}$.

Finally, suppose G that contains a parabolic. Then all infinite order elements of G are parabolic with unique fixed point β. Therefore, they fix a Busemann function at β. By the same argument as above, any elliptic element in G also fixes the Busemann function. □

In [KN04] Karlsson-Noskov studied isometry groups of visibility spaces with nonempty $X_G(\infty)$. To state their result we call a group G *no-F_2-connected* if G is generated by a set S of infinite order elements such that the following graph is connected: the vertices are elements of S, and two vertices $s_1, s_2 \in S$ are joined by an edge if and only if the subgroup generated by s_1, s_2 does not contain a nonabelian free subgroup.

By [KN04, Theorem 2] and Proposition 4.1 if G is a no-F_2-connected, not virtually-cyclic, discrete isometry group of a visibility manifold, then G is grounded, and it fixes a Busemann function.

The class of no-F_2-connected groups trivially contains each torsion-free group without free nonabelian subgroups. Also the product of any two nontrivial, torsion-free groups is no-F_2-connected (here S is the set of all elements in the product for which one of the coordinates equals the identity). More examples of no-F_2-connected groups can be found in [KN04, Section 8].

We conclude that if X visibility, then Part (3) of Theorem 3.5 can be improved to "G is amenable and not virtually cyclic" or by "G does not contains a non-abelian free subgroup and is generated by the set of its infinite order elements", and similarly, Part (5) can be improved to "an normal abelian subgroup of G is not cyclic".

5. Manifolds with $\mathbb{R}$-factors

In this section we discuss when an open manifold is homeomorphic to the product of $\mathbb{R}$ and another manifold. For background on ends of manifolds we refer to [Sie65, Geo08, GT03].

Guilbault showed in [Gui07] that if W is an open manifold that is homotopy equivalent to a finite complex, then $W \times \mathbb{R}$ has stable fundamental group at infinity and zero Siebenmann's end obstruction, so that if $\dim(W) \geq 5$, then $W \times \mathbb{R}$ is diffeomorphic to the interior of a compact manifold. If the complex has codimension ≥ 3 in $W \times \mathbb{R}$, one can say more:

Theorem 5.1. *Let W be an open m-manifold that is homotopy equivalent to a finite complex of dimension $k \leq m-2$. If $m \neq 3$, then $W \times \mathbb{R}$ is diffeomorphic to the interior of a regular neighborhood of k-dimensional subcomplex.*

Proof. The case $m = 2$ follows from the classification of open surfaces [Ric63], which implies that W is diffeomorphic to $\mathbb{R}^2$. Suppose $m \geq 4$ so that $V := W \times \mathbb{R}$ has dimension ≥ 5. Stalling's theorem [Sta] allows to embed any codimension ≥ 3 finite complex up to homotopy type, so let K be a finite k-dimensional subcomplex onto which V deformation retracts. Let R be a regular neighborhood of K. The inclusions $\partial R \to R \to V$ induce π_1-isomorphisms as K has codimension ≤ 3. Excision in the universal cover as in the proof of [Sie69, Theorem 2.1] implies that $U := V \setminus \mathrm{Int}(R)$ deformation retracts onto $\partial U = \partial R$.

The manifolds V and W are one-ended as their homology groups (with twisted coefficients) vanish in degrees $\geq m-2$ [Sie69, Proposition 1.2]. Denote their ends by ε_V, ε_W. By [Gui07, Propositions 3.1, 3.2] the end ε_V has arbitrary small finitely dominated 0-neighborhoods and its fundamental group is stable. Let $\pi_1(\infty_V)$ denote the inverse limit of the fundamental groups of the 0-neighborhoods; this group is finitely presented being a retract of the (finitely presented) fundamental group of a 0-neighborhood.

As R is compact, the above discussion of ε_{V} applies to the end ε_{U} of U, and in particular, $\pi_1(\infty_{U})$ and $\pi_1(\infty_{V})$ are isomorphic via the inclusion. If we show that the canonical map $e_{U}\colon \pi_1(\infty_{U}) \to \pi_1(U)$ is an isomorphism, then Siebenmann's open collar theorem [Sie69, Theorem 1.6] would imply that U is diffeomorphic to $\partial U \times [0,1)$, which would prove the theorem. Since the inclusion $U \to V$ induces a π_1-isomorphism, e_{U} is essentially the canonical map $e_{V}\colon \pi_1(\infty_{V}) \to \pi_1(V)$.

Next we note that e_{U} is onto and has perfect kernel. To see that multiply U by a circle to make the dimension ≥ 6. By Siebenmann's product formula for the end obstruction [Sie65, Theorem 7.5], $S^1 \times U$ is the interior of a compact manifold, which is a one-sided h-cobordism that deformation retracts onto the boundary component of the form $S^1 \times \partial U$. The other boundary component B satisfies $\pi_1(B) \cong \pi_1(S^1) \times \pi_1(\infty_{U})$. Restricting the retraction to B gives a map $B \to S^1 \times \partial U$, which is a π_1-surjection with perfect kernel by a standard argument involving Poincaré duality in the universal cover as e.g. in [GT03, Theorem 2.5]. Under the obvious identifications this π_1-surjection corresponds to the product of e_{U} and the identity of $\pi_1(S^1)$, so e_{U} is surjective with perfect kernel.

On the other hand, it follows from [Gui07, Propositions 3.5] that $\pi_1(\infty_{V})$ is isomorphic to a (possibly trivial) amalgamated product of the form $\pi_1(W) *_{\Gamma} \pi_1(W)$, and moreover, under the isomorphism e_{V} is identified with the standard projection $\pi_1(W) *_{\Gamma} \pi_1(W) \to \pi_1(W)$ that extends the identity maps on each factor. The kernel of this is a free group. (Indeed, the kernel is a normal subgroup that intersects the factors trivially, and hence intersects trivially each of their conjugates. Thus the kernel acts freely on the Bass-Serre tree for the amalgamated product, and hence it is a free group). Since a free perfect group must be trivial, e_{U} is an isomorphism, which finishes the proof. □

Remark 5.2. If $m = 3$ and $\pi_1(W)$ is trivial or $\mathbb{Z}$, which are the fundamental groups of $m-2$-complexes for which the topological surgery is known to work, then the same proof, based on the open collar theorem of [Gui92], shows that $W \times \mathbb{R}$ is homeomorphic to the interior of a regular neighborhood of K, i.e. D^4, or $D^3 \times S^1$, respectively.

Remark 5.3. The product of $\mathbb{R}$ with any aspherical $(n-1)$-manifold is covered by $\mathbb{R}^n$. In fact, the references below imply that the product of any two open contractible manifolds is diffeomorphic to $\mathbb{R}^n$ (provided their dimensions are positive and add up to n). This is due to Stallings [Sta62] if $n \geq 5$. The case of the product of an open contractible 3-manifold with $\mathbb{R}$ is due to Luft [Luf87, Theorem 5] and Perelman's solution of Poincaré conjecture. Finally, any contractible open 2-manifold is diffeomorphic to $\mathbb{R}^2$ by the classification of surfaces [Ric63]. These results only give a PL homeomorhism onto

$\mathbb{R}^n$, but a smooth structure on $\mathbb{R}^n$ is uniquely determined by the PL structure it induces [Mun60, Corollary 6.6].

Corollary 5.4. *Let $F \to E \to B$ be a fiber bundle whose base B is a closed manifold with $\pi_2(B) = 0$, and fiber F is a compact contractible m-manifold. If Int(E) is homeomorphic to $W \times \mathbb{R}$ for a manifold W, then Int(F) is homeomorphic to $\mathbb{R}^m$.*

Proof. By the h-cobordism theorem it suffices to show that ∂F is simply-connected, so we may assume $m \geq 3$. By Theorem 5.1 and Remark 5.2 the bundle projection restricted to ∂E induces a π_1-isomorphism, so since $\pi_2(B) = 0$, the homotopy sequence of the fibration $\partial F \to \partial E \to B$ implies $\pi_1(\partial F) = 0$. ☐

Corollary 5.5. *Let B be a closed manifold and C be an open aspherical m-manifold such that $B \times C$ are homeomorphic to the product of $\mathbb{R}$ and an open manifold. Then the universal cover of C is homeomorphic to $\mathbb{R}^m$.*

Proof. We can assume $\dim(C) \geq 3$, and that C is contractible, because the property of being homeomorphic to the product of $\mathbb{R}$ and an open manifold in inherited by any cover. Let W be an open manifold such that $B \times C$ is homeomorphic to $W \times \mathbb{R}$. By [Gui07, Propositions 3.1-3.2] $W \times \mathbb{R}$, and hence $B \times C$, has only one end at which π_1 is stable, and $\pi_1(\infty)$ is finitely presented. Hence the same holds for the end of C. It follows that $\pi_1(\infty)$ is the product of $\pi_1(B) \times \pi_1(\infty_C)$ and the projection on the first factor can be identified with the canonical map $\pi_1(\infty) \to \pi_1(B \times C)$, which is an isomorphism by Theorem 5.1 and Remark 5.2. Thus $\pi_1(\infty_C)$ is trivial, and hence C is homeomorphic to $\mathbb{R}^n$ by [Sta62, HP70, Gui92]. ☐

Proof of Corollary 1.5. If W is homeomorphic to $\mathbb{R}^n$, then it is diffeomorphic to $\mathbb{R}^n$ when $n \neq 4$, and $W \times S^1$ is diffeomorphic to $\mathbb{R}^4 \times S^1$ [Sie69, Theorem 2.2]. Of course, $\mathbb{R}^n \times S^1$ admits a metric of $K \leq 0$. Conversely, suppose $W \times S^1$ admits a metric of $K \leq 0$, and write it as X/G where G is infinite cyclic. The generator of G is either axial or parabolic. In the former case X/G is a vector bundle over S^1 as the normal exponential map to a totally geodesic submanifold of X is a diffeomorphism, and the bundle is trivial because $W \times S^1$ is orientable. In the latter case G fixes a Busemann function, so X/G is the product of $\mathbb{R}$ and an open manifold, and the claim follows from Corollary 5.5. ☐

6. Manifolds without $\mathbb{R}$-factors

The purpose of this section is to give a method of constructing $\mathbb{R}^n$-covered manifold that are not homeomorphic to $\mathbb{R}^{n-1} \times \mathbb{R}$ covered manifolds.

Theorem 6.1. *Let V be an open manifold that is covered by $\mathbb{R}^n$ for $n \geq 5$, and is homotopy equivalent to a finite complex of dimension $\leq n-3$. If $G := \pi_1(V)$ is nontrivial, there are countably many open $K(G,1)$ manifolds that are covered by $\mathbb{R}^n$ but are not homeomorphic to manifolds covered by $\mathbb{R} \times \mathbb{R}^{n-1}$.*

Proof. By Stalling's theorem [Sta] allows to embed any codimension ≥ 3 finite complex up to homotopy type, so let $N \subset V$ be a regular neighborhood of that subcomplex. Since the subcomplex has codimension ≤ 3, we know that ∂N is connected and the inclusions $\partial N \to N \to V$ induce π_1-isomorphisms. The universal cover of $\mathrm{Int}(N)$ is diffeomorphic to $\mathbb{R}^n$ because it is a regular neighborhood of a codimension ≥ 3 subcomplex, which easily implies that it is simply-connected at infinity, as mentioned e.g. in the statement of [Sie69, Theorem 2.1].

Thus there is an embedded circle in B that is homotopically nontrivial in N. By replacing it with a embedded circle homotopic to the square of the original one, we may assume its normal bundle is trivial (and it is still homotopically nontrivial in $\pi_1(N)$ since this group is torsion-free). Denote the curve by α, and its closed tubular neighborhood by T_α.

Let C be any compact contractible $(n-1)$-manifold bounding a non-simply-connected homology sphere whose fundamental group is freely indecomposable see e.g. Appendix B, or [CK65, Gla67]. Fix a closed coordinate disk $\Delta \subset \partial C$, attach $C \times S^1$ to N by identifying $\Delta \times S^1$ with T_α, and denote the resulting compact manifold with boundary by $N_{\alpha,C}$ (the choices involved in this identification will be irrelevant for our purposes).

The fundamental group of the boundary of $N_{\alpha,C}$ is the amalgamated product of $\pi_1(\partial N)$ and $\pi_1(\partial C) \times \mathbb{Z}$ along the infinite cyclic subgroup generated by the homotopy class of α. Taking the quotient by the normal closure of $\pi_1(\partial N)$ we see that the amalgamated product surjects onto $\pi_1(\partial C)$, so its rank is at least as large as the rank of $\pi_1(\partial C)$.

By taking boundary connected sums of C with itself one gets a sequence $\{C_i\}_{i \in \mathbb{N}}$ of compact contractible manifolds such that C_i bounds the connected sum Σ_i of i copies of the homology sphere $\Sigma := \partial C$. Grushko's theorem implies $\mathrm{rank}(\pi_1(\Sigma_i)) = i\, \mathrm{rank}(\pi_1(\Sigma))$, so $V_i := \mathrm{Int}(N \#_{\alpha,C_i})$ are pairwise non-homeomorphic because their fundamental groups of infinity are pairwise non-isomorphic. Deformation retraction $C_i \to \Delta$ extends to a deformation retraction of $N_{\alpha,C_i} \to N$, so V_i is homotopy equivalent to V.

Next we show that V_i is covered by $\mathbb{R}^n$. Denote the universal cover of N and V by $\widetilde{N}$ and $\widetilde{V}$, respectively. Since α is homotopically nontrivial, it unravels in $\widetilde{N}$ to a proper embedding $\mathbb{R} \to \partial \widetilde{N}$. Let T be the union of all lifts of T_α to $\partial \widetilde{N}$. Each component of T is an open disk, and these disks have disjoint closures. Then $\widetilde{V} \cup T$ is a noncompact manifold with boundary T. Note that $\widetilde{V} \cup T$ can be compactified to D^n where T becomes the union of a countable collection of round open disks with pairwise disjoint closures, namely, attach a collar to T and invoke Cantrell-Stalling's hyperplane linearization theorem (see [CKS, Corollary 9.3] for a new proof and references to original proofs). On the other hand, the universal cover of $S^1 \times (\mathrm{Int}(\Delta) \cup \mathrm{Int}(C_i))$ is a noncompact manifold with boundary whose interior is diffeomorphic to $\mathbb{R}^n$ [Sta62] and whose boundary is an open disk, so it also compactifies to D^n so that the disk $\mathbb{R}\times\mathrm{Int}(\Delta)$ becomes a round disk. It follows that attaching each lift of $S^1 \times (\mathrm{Int}(\Delta) \cup \mathrm{Int}(C_i))$ to $\tilde{V} \cup T$ simply amounts to attaching an open collar to each component of T, and hence the result has interior diffeomorphic to $\widetilde{V}$, which is $\mathbb{R}^n$.

It remains to show that V_i is not homeomorphic to a manifold covered by $\mathbb{R} \times \mathbb{R}^{n-1}$. If it is, then the same is true for $T^r \times V_i$ where $r \geq 1$ to ensure that $T^r \times V_i$ has dimension ≥ 6. Hence Theorem 5.1 applies and the map $\pi_1(\infty) \to \pi_1(S^1 \times V_i)$ is an isomorphism. Hence the same is true for the boundary inclusion into $S^1 \times N_{\alpha,C_i}$, which on the other hand is not injective because any homotopically nontrivial loop in Σ_i is null-homotopic in C_i. This contradiction completes the proof. □

We are able to replace the conclusion “countably many” in Theorem 6.1 by ”a continuum of” only under the technical assumption introduced below (in which *splits* stands for *splits as an amalgamated product or an HNN-extension*).

Let $\{\Sigma_i^3\}_{i\in\mathbb{N}}$ be an infinite family of aspherical homology 3-spheres Σ_i that bound contractible manifolds and such that $\pi_1(\Sigma_i^3 \times S^1)$ are pairwise non-isomorphic (see Appendix B for examples of such families). Consider the sequence of groups $\mathfrak{S} := \{\pi_1(\Sigma_i^3 \times S^1)\}_{i\in\mathbb{N}}$. We say that a group G is *no-*$\mathfrak{S}$ if **one** of the following conditions (1), (2), (3) holds:

1. no group in $\mathfrak{S}$ can be embedded into G.
2. G splits as a (possibly infinite and possibly consisting of a single vertex) graph of groups such that
   - (a) all edge groups have cohomological dimension ≤ 2,
   - (b) no vertex group is isomorphic to a group in $\mathfrak{S}$.
   - (c) no vertex group splits over a trivial or an infinite cyclic subgroup.

I believe there should exist a group of type F that is not no-$\mathfrak{S}$ for every $\mathfrak{S}$. Here we are more interested in groups that are no-$\mathfrak{S}$ for some $\mathfrak{S}$.

Example 6.2. The fundamental group G of any closed aspherical n-manifold (or more generally, any Poincaré duality group) is no-$\mathfrak{S}$ for any $\mathfrak{S}$. This is trivial for $n \leq 3$ as $\mathrm{cd}(G) \leq 3$ so (1) holds, and if $n \geq 3$, than a Mayer-Vietoris sequence in group cohomology shows that G does not split as an amalgamated product over a subgroup of cohomological dimension ≤ 1, so (2) holds.

Example 6.3. Appendix B contains an infinite family of Σ_i's that are modelled on $\widetilde{SL}_2(\mathbb{R})$ geometry and are found by Casson-Harer. For this family the groups $\mathbb{Z}^3$ and $\mathbb{Z}*\mathbb{Z}$ embed into every group in $\mathfrak{S}$. Thus if $\mathbb{Z}^3$ or $\mathbb{Z}*\mathbb{Z}$ does not embed into G, then G is no-$\mathfrak{S}$ by (1).

Remark 6.4. If G splits as a finite graph of groups that satisfies (a) and (c), then it can be arranged to satisfy (b) by removing the (finitely many) vertex groups from $\mathfrak{S}$. Similarly, if G violates (1) for every $\mathfrak{S}$, then G contains all but finitely many groups of the form $\pi_1(\Sigma^3 \times S^1)$ where Σ^3 is an aspherical homology 3-sphere that bounds a contractible manifolds.

Theorem 6.5. *Let V be an open aspherical n-manifold that is homotopy equivalent to a finite complex of dimension $\leq n-3$, where $n \geq 5$ and $\pi_1(V)$ is no-$\mathfrak{S}$ for some $\mathfrak{S}$. Then the tangential homotopy type of V contains a continuum of manifolds covered by $\mathbb{R}^n$, and not homeomorphic to manifolds covered by $\mathbb{R}\times\mathbb{R}^{n-1}$.*

Proof. As in the proof of Theorem 6.1 we can change V within its tangential homotopy type so that assume V is the interior of a regular neighborhood N of a codimension ≥ 3 finite subcomplex. Set $B := \partial N$, so that $\pi_1(B) \to \pi_1(N) \cong \pi_1(V)$ is an isomorphism induced by the inclusion.

Results of Kervaire, recalled in Appendix B enure the existence of an infinite sequence of compact contractible $(n-1)$-manifolds $\{C_i\}_{i\in\mathbb{N}}$ such that $\pi_1(\partial C_i) \cong \pi_1(\Sigma_i^3) \in \mathfrak{S}$. Set $\Sigma_i := \partial C_i$.

Pick two disjoint closed coordinate disks Δ_{i-}, Δ_{i+} in Σ_i. Given a subset $\alpha \subseteq \mathbb{N}$, let C_α be a boundary connected sum of C_i's with indices in α defined as follows: if $\alpha = \{i_1, \dots, i_k, \dots\}$ with the ordering induced from $\mathbb{N}$, then we attach C_{i_k} to $C_{i_{k+1}}$ by identifying Δ_{i_k+} and $\Delta_{i_{k+1}-}$ for all k.

Pick a nontrivial element in $\pi_1(B)$. By replacing the element with its square, which we denote s, we can assume that it can be represented by an embedded circle in B with trivial normal bundle. Attach N to $C_\alpha \times S^1$ by identifying $\Delta_{i_1-} \times S^1$ with a closed tubular neighborhood of the above embedded circle in B via an arbitrary homeomorphism, denote the result by N_α, and set $V_\alpha := \mathrm{Int}(N_\alpha)$. Since N_α deformation retracts to N, the manifolds V_α and V are tangentially homotopy equivalent. The corresponding part of the proof of Theorem 6.1 implies that V_α is covered by $\mathbb{R}^n$ but is not homeomorphic to a manifold covered by $\mathbb{R}\times\mathbb{R}^{n-1}$. It remains to show that $\{V_\alpha\}$ fall into a continuum of homeomorphism types.

We next construct a nested sequence $\{W_k\}$ of codimension zero submanifolds that are neighborhood of infinity in $\mathrm{Int}(C_\alpha)$ such that $\mathrm{Int}(W_k) \supset W_{k+1}$, the boundary ∂W_k is compact, and $\bigcap_{k\geq 1} W_k = \emptyset$. (Any sequence of neighborhoods that satisfy the last property is called *cofinal*). Imagine a worm that eats the inside of C_{i_1} except for a collar neighborhood of Σ_{i_1}, which gives W_1. For the second helping the worm eats some more of $\mathrm{Int}(C_{i_1})$ leaving a smaller collar neighborhood of Σ_{i_1}, and then eats a tunnel into C_{i_2} through Δ_{i_2-}, by leaving a collar neighborhood of $\partial\Delta_{i_2-}$ in $\Delta_{i_2}-$, and then eats the inside of C_{i_2} except for a collar neighborhood of Σ_{i_2}, which results in W_2. The process is repeated so that the worm gets into C_{i_k} on the step k, which yields W_k. Then $\pi_1(W_1) \cong \pi_1(\Sigma_{i_1})$, and in general $\pi_1(W_k) \cong \pi_1(\Sigma_{i_1}) * \cdots * \pi_1(\Sigma_{i_k})$.

Now $\{W_k \times S^1\}$ is a sequence of neighborhoods of infinity in $\mathrm{Int}(C_\alpha) \times S^1$. Let $\{O_k\}$ be a nested cofinal sequence of neighborhoods of infinity of $\mathrm{Int}(N)$ such that $O_k \bigcup B$ is a collar neighborhood of B. We let the worm enlarge $O_k \cup (W_k \times S^1)$ by eating through the interior of $\Delta_{i_1-} \times S^1$, except for a collar of $\partial\Delta_{i_1-} \times S^1$, until it connects to O_k; we denote the result by U_k. Thus $\{U_k\}$ is a nested, cofinal sequence of neighborhood of infinity $\{U_k\}_{k\geq 1}$ in $V_\alpha = \mathrm{Int}(N_\alpha)$ such that $\mathrm{Int}(U_k) \supset U_{k+1}$ and the boundary ∂U_k is compact. By van Kampen's theorem $\pi_1(U_k)$ is the amalgamated product of $\pi_1(W_k \times S^1)$ and $\pi_1(B)$ along the infinite cyclic subgroup generated by s.

Given basepoints $p_k \in U_k$ and paths r_k in U_k connecting p_k to p_{k+1}, we get an inverse sequence of homomorphisms $\{\rho_k\colon\ \pi_1(U_k, p_k) \longleftarrow \pi_1(U_{k+1}, p_{k+1})\}$ induced by the inclusion $U_{k+1} \subset U_k$ followed by the basepoint change conjugation via r_k's. Each ρ_k has a section defined as the composition of homomorphisms induced by the following operations: deformatation retract $\bigcup_{m>k} C_{i_m} \times S^1$ onto $\Delta_{i_k+} \times S^1$ inside U_k, intersect the result with U_{k+1}, include the intersection into U_{k+1}, and change the base point via r_k. Except for including into U_{k+1}, these operations induce fundamental group isomorphisms.

Two inverse sequence of homomorphisms $\{A_l \longleftarrow A_{l+1}\}$, $\{B_m \longleftarrow B_{m+1}\}$ are *pro-equivalent* if after passing to subsequences there is a commutative diagram

$$\begin{tikzcd}
A_{l_1} & & A_{l_2} \arrow[ll] \arrow[dl] & & A_{l_3} \arrow[ll] \arrow[dl] & & \cdots \arrow[ll] \arrow[dl] \\
& B_{m_1} \arrow[ul] & & B_{m_2} \arrow[ll] \arrow[ul] & & B_{m_3} \arrow[ll] \arrow[ul] & & \cdots \arrow[ll]
\end{tikzcd}$$

in which the horizontal arrows are the composites of the given homomorphisms (see e.g. [GG, 2.1] for details). A sequence of homomorphisms is *semistable* if it is pro-equivalent to a sequence of surjective homomorphisms.

Since ρ_k has a section, it is surjective, so the sequence $\{\rho_k\}$ is semistable, which implies (see e.g. [GG, 2.2]) that its pro-equivalence class is independent of any of the choices of U_k, p_k, r_k, and hence it depends only on the homeomorphism type of $\mathrm{Int}(N_\alpha)$. If we set $G_k^\alpha := \pi_1(U_k, p_k)$, and identify G_k^α with a subgroup of G_{k+1}^α via the above section, then ρ_k becomes a retraction $G_k^\alpha \longleftarrow G_{k+1}^\alpha$.

For the rest of the proof we think of G_k^α as an amalgamated product of subgroups isomorphic to $\pi_1(B), \pi_1(\Sigma_{i_1} \times S^1), \dots, \pi_1(\Sigma_{i_k} \times S^1)$ along $\langle s \rangle$ and refer to these subgroups as *factors*. Note that G_k^α is the subgroup of G_{k+1}^α generated by the above factors, and the kernel of the retraction $G_k^\alpha \longleftarrow G_{k+1}^\alpha$ equals the normal closure of the subgroup $\pi_1(\Sigma_{i_{k+1}}) \times \{1\}$ of the $\pi_1(\Sigma_{i_{k+1}}) \times \mathbb{Z}$-factor, which is mapped to G_k^α by projection onto $\mathbb{Z}$, identified with $\langle s \rangle$.

To get a continuum of homeomorphism types of $\mathrm{Int}(N_\alpha)$'s it suffices to show that that the pro-equivalence of $\{G_k^\beta \longleftarrow G_{k+1}^\beta\}$ and $\{G_k^\gamma \longleftarrow G_{k+1}^\gamma\}$ implies equality of the subsets β, γ, and by symmetry we only need to show that any $i_k \in \beta$ also lies in γ. Look at a portion of the commutative diagram from the definition of pro-equivalence:

$$\begin{tikzcd}
G_{l_1}^\beta & & G_{l_2}^\beta \arrow[ll, "q_1"'] \arrow[dl] & & G_{l_3}^\beta \arrow[ll, "q_2"'] \arrow[dl, "\psi"] \\
& G_{m_1}^\gamma \arrow[ul] & & G_{m_2}^\gamma \arrow[ll, "p"'] \arrow[ul, "\phi"] &
\end{tikzcd}$$

with $l_2 > l_1 > i_k$ and $m_2 > m_1 > i_k$. With the above identifications q_1 and q_2 both restrict to the identity on G_k^β. Commutativity implies that $\psi \circ p$ and ψ restricted to G_k^β is injective, and hence so is ϕ restricted to $\psi(G_k^\beta)$.

Let $Q \le G_{l_3}^\beta$ denote the subgroup corresponding to the $\pi_1(\Sigma_{i_k} \times S^1)$-factor. Note that Q does not split over a trivial or an infinite cyclic subgroup because $\Sigma_{i_k}^3 \times S^1$ is a closed aspherical 4-manifold, and the fundamental group of a closed aspherical manifold of dimension > 2 cannot have such splittings.

Since $G_{m_2}^\gamma$ is an amalgamated product over $\langle s \rangle$, the group $\psi(Q)$ lies in a conjugate, denoted R, of one of the factors of $G_{m_2}^\gamma$. The map p restricted to a conjugate of a factor is either injective or is a projection onto $\langle s \rangle$. The latter is impossible for $p|_R$ since the image contains $p(\psi(Q)) \cong Q$. It follows that ϕ restricted to R is injective. Now R is isomorphic either to $\pi_1(B)$ or to a group in $\mathfrak{S}$. Let us consider the former case, and use that $\pi_1(B) \cong \pi_1(V)$ is no-$\mathfrak{S}$. As $\psi(Q) \le R$, the assumption (1) gives a contradiction. The assumption (2) implies that $\psi(Q)$ lies in a vertex group R_0 in a graph of groups decomposition of $\pi_1(N)$ as in (2) such that R_0 does not split over a trivial or an infinite cyclic subgroup.

Then $\phi(R_0)$ lies in a conjugate, denoted S, of one of the factors of $G_{l_2}^\beta$, which implies $S \supset \phi(\psi(Q)) = Q$. It follows that $S = Q$ for otherwise S fixes a vertex (in the Bass-Serre tree of the amalagamated product decomposition of G_l^β) that is distinct from the vertex fixed by Q, so Q would fix the path joining the two vertices, but edge stabilizers are cyclic and Q is not. Thus

$Q = \phi(\psi(Q)) \subseteq \phi(R_0) \subseteq S = Q$, so injectivity of ϕ on R_0 implies $R_0 = \phi(Q)$, which contradicts (b).

Finally, if R is isomorphic to a group in $\mathfrak{S}$, then R also does not split over a trivial or an infinite cyclic subgroup, so the argument of the previous paragraph gives $R = \psi(Q)$, which means that a group in $\mathfrak{S}$ is among the factors of $G^{\gamma}_{m_2}$ other than the $\pi_1(B)$-factor, i.e. i_k lies in γ. □

Appendix A. Finiteness properties for groups

Our main results apply to groups of type *F* or groups of finite cohomological dimension. This appendix contains references justifying that various groups constructed in Sections 2-3 have these properties (provided they are torsion-free). We refer to [Bie81, Bro94] for properties of cohomological and geometric dimensions of a group, of which we mostly use the former, denoted cd. Finiteness conditions on groups, such as *F* and *FP*, are reviewed in [Dav08, Appendix F]. The following facts are known:

1. Finiteness of cohomological dimension in inherited by extensions and amalgamated products [Bro94, Proposition VIII.2.4].
2. The property of having type *F* is (clearly) inherited by amalgamated products, i.e. if $G_1 *_A G_2$ has type F when so do G_1, G_2, and A.
3. The property of having type *F* or *FP* is inherited by extensions. (Combine [Bro94, Exercise VIII.6.8] with the fact that finite presentability of a group inherited by extensions [Rob96, 2.4.4])
4. The fundamental group of an aspherical n-manifold has cohomological dimension $\leq n$ with equality precisely when the manifold is closed [Bro94, VIII.8.1]. More generally, an infinite index subgroup of the n-dimensional Poincaré duality group has cohomological dimension $< n$ [Str77].
5. G is a countable group of finite cohomological dimension if and only if there is a $K(G,1)$ manifold whose universal cover is a Euclidean space. (Find a finite-dimensional $K(G,1)$ [Bro94, VIII.7.1], make it a locally finite and of the same dimension [Whi49, Theorem 13], use simplicial approximation to make the space polyhedral, embed it into a Euclidean space [Spa81, Theorem 3.2.9], take a regular neighborhood, and note that the universal cover of any codimension ≥ 3 regular neighborhood is simply-connected at infinity, as mentioned e.g. in the statement of [Sie69, Theorem 2.1]. Alternatively, one can take the product of the regular neighborhood with $\mathbb{R}$, see Remark 5.3.)

Appendix B. Homology spheres bounding contractible manifolds

Results of Section 6 require a certain infinite family of compact contractible manifolds whose construction is reviewed below.

The problem which homology 3-spheres bound contractible manifolds is not well understood. (Freedman famously showed that all homology 3-spheres bound topological contractible 4-manifolds, but in this paper we only consider smooth manifolds). An infinite family of homology 3-spheres that bound contractible manifolds was found by Casson-Harer [CH81] among Brieskorn spheres $\Sigma(p,q,r)$ with $\frac{1}{p}+\frac{1}{q}+\frac{1}{r}<1$, which are closed aspherical 3-manifolds modelled on $\widetilde{SL}_2(\mathbb{R})$ geometry [Mil75]. For example, Casson-Harer's list contains the family $\Sigma(p,p+1,p+2)$ where p is any odd integer ≥ 3. The groups $\pi_1(\Sigma(p,p{+}1,p{+}2)\times S^1)$ are pairwise non-isomorphic because their quotients by the centers is the fundamental groups of 2-orbifolds with Euler characteristics $\frac{1}{p}+\frac{1}{p+1}+\frac{1}{p+2}-1$, which are all distinct.

One can also construct an infinite family of hyperbolic homology 3-spheres that bound contractible manifolds, see the MathOverflow answers [AM] for details, but the above Casson-Harer's family is enough for our purposes.

Kervaire shows in [Ker69, Theorems 1 and 3] that for each $m>3$ the fundamental group of any homology 3-sphere is also the fundamental group of a homology m-sphere that bounds a contractible manifold. (Even though [Ker69, Theorems 1] is not stated for $m=4$, it still applies in our case because as Kervaire remarks on [Ker69, pp. 67-68] the fundamental group of any homology 3-sphere has a presentations with an equal number of relations and generators).

References

[AJ76] L. Auslander and F. E. A. Johnson, *On a conjecture of C. T. C. Wall*, J. London Math. Soc. (2) **14** (1976), no. 2, 331–332.

[AM] I. Agol and B. Martelli, *Irreducible homology 3-spheres that bound smooth contractible manifolds*, http://mathoverflow.net/questions/104451/irreducible-homology-3-spheres-that-bound-smooth-contractible-manifolds.

[AS82] R. C. Alperin and P. B. Shalen, *Linear groups of finite cohomological dimension*, Invent. Math. **66** (1982), no. 1, 89–98.

[Bau93] G. Baumslag, *Topics in combinatorial group theory*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 1993.

[BB76] G. Baumslag and R. Bieri, *Constructable solvable groups*, Math. Z. **151** (1976), no. 3, 249–257.

[BdlHV08] B. Bekka, P. de la Harpe, and A. Valette, *Kazhdan's property (T)*, New Mathematical Monographs, vol. 11, Cambridge University Press, Cambridge, 2008.

[BF02] M. Bestvina and M. Feighn, *Proper actions of lattices on contractible manifolds*, Invent. Math. **150** (2002), no. 2, 237–256.

[BGS85] W. Ballmann, M. Gromov, and V. Schroeder, *Manifolds of nonpositive curvature*, Progress in Mathematics, vol. 61, Birkhäuser Boston Inc., Boston, MA, 1985.

[BH99] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 319, Springer-Verlag, Berlin, 1999.

[Bie81] R. Bieri, *Homological dimension of discrete groups*, second ed., Queen Mary College Mathematical Notes, Queen Mary College Department of Pure Mathematics, London, 1981.

[BKK02] M. Bestvina, M. Kapovich, and B. Kleiner, *Van Kampen's embedding obstruction for discrete groups*, Invent. Math. **150** (2002), no. 2, 219–235.

[Bre72] G. E. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972, Pure and Applied Mathematics, Vol. 46.

[Bro94] K. S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York, 1994, Corrected reprint of the 1982 original.

[BS73] A. Borel and J.-P. Serre, *Corners and arithmetic groups*, Comment. Math. Helv. **48** (1973), 436–491, Avec un appendice: Arrondissement des variétés à coins, par A. Douady et L. Hérault.

[BS87] M. Burger and V. Schroeder, *Amenable groups and stabilizers of measures on the boundary of a Hadamard manifold*, Math. Ann. **276** (1987), no. 3, 505–514.

[CH81] A. J. Casson and J. L. Harer, *Some homology lens spaces which bound rational homology balls*, Pacific J. Math. **96** (1981), no. 1, 23–36.

[CK65] M. L. Curtis and K. W. Kwun, *Infinite sums of manifolds*, Topology **3** (1965), 31–42.

[CKS] J. S. Calcut, H. C. King, and L. C. Siebenmann, *Connected sum at infinity and cantrell-stallings hyperplane unknotting*, arXiv:1010.2707v1.

[CL10] P.-E. Caprace and A. Lytchak, *At infinity of finite-dimensional CAT(0) spaces*, Math. Ann. **346** (2010), no. 1, 1–21.

[CM09] P.-E. Caprace and N. Monod, *Isometry groups of non-positively curved spaces: structure theory*, J. Topol. **2** (2009), no. 4, 661–700.

[Dav08] M. W. Davis, *The geometry and topology of Coxeter groups*, London Mathematical Society Monographs Series, vol. 32, Princeton University Press, Princeton, NJ, 2008.

[Dek96] K. Dekimpe, *Almost-Bieberbach groups: affine and polynomial structures*, Lecture Notes in Mathematics, vol. 1639, Springer-Verlag, Berlin, 1996.

[Des06] Z. Despotovic, *Action dimension of mapping class groups*, ProQuest LLC, Ann Arbor, MI, 2006, Thesis (Ph.D.)–The University of Utah.

[DIKL95] K. Dekimpe, P. Igodt, S. Kim, and K.B. Lee, *Affine structures for closed* 3*-dimensional manifolds with nil-geometry*, Quart. J. Math. Oxford Ser. (2) **46** (1995), no. 182, 141–167.

[DJL12] M. Davis, T. Januszkiewicz, and J.-F. Lafont, 4*-dimensional locally* $\mathrm{CAT}(0)$*-manifolds with no Riemannian smoothings*, Duke Math. J. **161** (2012), no. 1, 1–28.

[Duc] B. Duchesne, *Superrigidity in infinite dimension and finite rank via harmonic maps*, arXiv:1206.1964v1.

[Ebe96] P. B. Eberlein, *Geometry of nonpositively curved manifolds*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1996.

[Geo08] R. Geoghegan, *Topological methods in group theory*, Graduate Texts in Mathematics, vol. 243, Springer, New York, 2008.

[GG] R. Geoghegan and C. R. Guilbault, *Topological properties of spaces admitting free group actions*, arXiv:1005.2766v2.

[Gla67] L. C. Glaser, *Uncountably many contractible open* 4*-manifolds*, Topology **6** (1967), 37–42.

[Gro03] M. Gromov, *Random walk in random groups*, Geom. Funct. Anal. **13** (2003), no. 1, 73–146.

[GT03] C. R. Guilbault and F. C. Tinsley, *Manifolds with non-stable fundamental groups at infinity. II*, Geom. Topol. **7** (2003), 255–286.

[Gui92] C. R. Guilbault, *An open collar theorem for 4-manifolds*, Trans. Amer. Math. Soc. **331** (1992), no. 1, 227–245.

[Gui07] ———, *Products of open manifolds with* $\mathbb{R}$, Fund. Math. **197** (2007), 197–214.

[Hae61] A. Haefliger, *Plongements différentiables de variétés dans variétés*, Comment. Math. Helv. **36** (1961), 47–82.

[HIH77] E. Heintze and H.-C. Im Hof, *Geometry of horospheres*, J. Differential Geom. **12** (1977), no. 4, 481–491 (1978).

[HL92] J. A. Hillman and P. A. Linnell, *Elementary amenable groups of finite Hirsch length are locally-finite by virtually-solvable*, J. Austral. Math. Soc. Ser. A **52** (1992), no. 2, 237–241.

[HP70] L. S. Husch and T. M. Price, *Finding a boundary for a 3-manifold*, Ann. of Math. (2) **91** (1970), 223–235.

[Ker69] M. A. Kervaire, *Smooth homology spheres and their fundamental groups*, Trans. Amer. Math. Soc. **144** (1969), 67–72.

[KM79] M. I. Kargapolov and Ju. I. Merzljakov, *Fundamentals of the theory of groups*, Graduate Texts in Mathematics, vol. 62, Springer-Verlag, New York, 1979, Translated from the second Russian edition by Robert G. Burns.

[KN04] A. Karlsson and G. A. Noskov, *Some groups having only elementary actions on metric spaces with hyperbolic boundaries*, Geom. Dedicata **104** (2004), 119–137.

[Kro86] P. H. Kropholler, *Cohomological dimension of soluble groups*, J. Pure Appl. Algebra **43** (1986), no. 3, 281–287.

[LS69] W. B. R. Lickorish and L. C. Siebenmann, *Regular neighbourhoods and the stable range*, Trans. Amer. Math. Soc. **139** (1969), 207–230.

[Luf87] E. Luft, *On contractible open 3-manifolds*, Aequationes Math. **34** (1987), no. 2-3, 231–239.

[Mal51] A. I. Mal′cev, *On some classes of infinite soluble groups*, Mat. Sbornik N.S. **28(70)** (1951), 567–588.

[Mes90] G. Mess, *Examples of Poincaré duality groups*, Proc. Amer. Math. Soc. **110** (1990), no. 4, 1145–1146.

[Mil75] J. Milnor, *On the 3-dimensional Brieskorn manifolds* $M(p,q,r)$, Knots, groups, and 3-manifolds (Papers dedicated to the memory of R. H. Fox), Princeton Univ. Press, Princeton, N. J., 1975, pp. 175–225. Ann. of Math. Studies, No. 84.

[Mun60] J. Munkres, *Obstructions to the smoothing of piecewise-differentiable homeomorphisms*, Ann. of Math. (2) **72** (1960), 521–554.

[NS11] A. Naor and L. Silberman, *Poincaré inequalities, embeddings, and wild groups*, Compos. Math. **147** (2011), no. 5, 1546–1572.

[Ont03] P. Ontaneda, *The double of a hyperbolic manifold and non-positively curved exotic PL structures*, Trans. Amer. Math. Soc. **355** (2003), no. 3, 935–965 (electronic).

[Ric63] I. Richards, *On the classification of noncompact surfaces*, Trans. Amer. Math. Soc. **106** (1963), 259–269.

[Rob96] D. J. S. Robinson, *A course in the theory of groups*, second ed., Graduate Texts in Mathematics, vol. 80, Springer-Verlag, New York, 1996.

[RS68] C. P. Rourke and B. J. Sanderson, *Block bundles. I*, Ann. of Math. (2) **87** (1968), 1–28. MR 0226645 (37 #2234a)

[Sap] M. Sapir, *Aspherical groups and manifolds with extreme properties*, arXiv:1103.3873v3.

[Sie65] L. C. Siebenmann, *The obstruction to finding a boundary for an open manifold of dimension greater than five*, ProQuest LLC, Ann Arbor, MI, 1965, Thesis (Ph.D.)–Princeton University.

[Sie69] ______, *On detecting open collars*, Trans. Amer. Math. Soc. **142** (1969), 201–227.

[Spa81] E. H. Spanier, *Algebraic topology*, Springer-Verlag, New York, 1981, Corrected reprint of the 1966 original.

[Sta] J. R. Stallings, *The embedding of homotopy types into manifolds*, 2000 version, http://math.berkeley.edu/~stall/embkloz.pdf.

[Sta62] J. Stallings, *The piecewise-linear structure of Euclidean space*, Proc. Cambridge Philos. Soc. **58** (1962), 481–488.

[Str77] R. Strebel, *A remark on subgroups of infinite index in Poincaré duality groups*, Comment. Math. Helv. **52** (1977), no. 3, 317–324.

[Szc85] A. Szczepański, *Euclidean space forms with the first Betti number equal to zero*, Quart. J. Math. Oxford Ser. (2) **36** (1985), no. 144, 489–493.

[Wal99] C. T. C. Wall, *Surgery on compact manifolds*, second ed., Mathematical Surveys and Monographs, vol. 69, American Mathematical Society, Providence, RI, 1999, Edited and with a foreword by A. A. Ranicki.

[Whi49] J. H. C. Whitehead, *Combinatorial homotopy. I*, Bull. Amer. Math. Soc. **55** (1949), 213–245.

[Wil00] B. Wilking, *On fundamental groups of manifolds of nonnegative curvature*, Differential Geom. Appl. **13** (2000), no. 2, 129–165.

[Yoo04] S. Y. Yoon, *A lower bound to the action dimension of a group*, Algebr. Geom. Topol. **4** (2004), 273–296 (electronic).

Igor Belegradek, School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332-0160

E-mail address: `ib@math.gatech.edu`